

UC Irvine

UC Irvine Previously Published Works

Title

Angelica keiskei, an emerging medicinal herb with various bioactive constituents and biological activities.

Permalink

<https://escholarship.org/uc/item/0qv5p87w>

Journal

Archives of pharmacal research, 40(6)

ISSN

0253-6269

Authors

Kil, Yun-Seo
Pham, Sally T
Seo, Eun Kyoung
et al.

Publication Date

2017-06-01

DOI

10.1007/s12272-017-0892-3

Peer reviewed

REVIEW

Angelica keiskei, an emerging medicinal herb with various bioactive constituents and biological activities

Yun-Seo Kil¹ · Sally T. Pham² · Eun Kyoung Seo¹ · Mahtab Jafari²

Received: 29 October 2016 / Accepted: 30 January 2017 / Published online: 24 April 2017
© The Pharmaceutical Society of Korea 2017

Abstract *Angelica keiskei* (Miq.) Koidz. (Umbelliferae) has traditionally been used to treat dysuria, dyschezia, and dysgalactia as well as to restore vitality. Recently, the aerial parts of *A. keiskei* have been consumed as a health food. Various flavonoids, coumarins, phenolics, acetylenes, sesquiterpene, diterpene, and triterpenes were identified as the constituents of *A. keiskei*. The crude extracts and pure constituents were proven to inhibit tumor growth and ameliorate inflammation, obesity, diabetics, hypertension, and ulcer. The extract also showed anti-thrombotic, anti-oxidative, anti-hyperlipidemic, anti-viral, and anti-bacterial activities. This valuable herb needs to be further studied and developed not only to treat these human diseases but also to improve human health. Currently *A. keiskei* is commercialized as a health food and additives in health drinks. This article presents a comprehensive review of *A. keiskei* and its potential place in the improvement of human health.

Keywords *Angelica keiskei* · Umbelliferae · Chemical constituent · Biological activity

Introduction

Angelica keiskei (Miq.) Koidz. is a hardy perennial, belonging to the family Umbelliferae. *A. keiskei* has traditionally been used as a diuretic, mild cathartic, tonic, and galactagogue (Kimura and Baba 2003). The herb is called ‘Myeong-II Yeob’ in Korea and ‘Ashitaba’ in Japan, both literally meaning ‘Tomorrow’s Leaf’ (Baba et al. 1998). Another common name for *A. keiskei* is ‘Shin-Sun Cho’, which means ‘a precious herb used by God’. *A. keiskei* is mainly distributed in Asian countries including Korea and Japan (Park 2013). The aerial parts are mainly used as a source of green vegetable juice as a daily health food and the young leaves of the aerial parts are used by parboiling or deep-frying (Imai and Imai 2008). The aerial parts are also used in tea, flour, wine, and cosmetics (Bao 2014; Su 2014; Ku et al. 2014; Liu et al. 2014). This review paper summarizes botanical characteristics, phytochemical research, as well as biological studies on the extracts and pure constituents of *A. keiskei*. The extracts have been reported to have anti-inflammatory, anti-obesity, anti-oxidative, anti-coagulant, anti-tumor, anti-mutagenic, anti-diabetic, anti-bacterial, and hepato-protective activities. The biological activities of each pure compounds are also separately summarized in this review.

Botanical characteristics of *A. keiskei*

Angelica keiskei has glossy leaves and flowers from summer to fall. The leaves are harvested from spring to fall before they lose the gloss. A yellow exudate is squeezed out of the stems when the leaves are harvested (Imai and Imai 2008). This yellow color makes it easy to distinguish *A. keiskei* from *A. japonica*, an exudate of which shows a

✉ Eun Kyoung Seo
yuny@ewha.ac.kr

✉ Mahtab Jafari
mjafari@uci.edu

¹ College of Pharmacy, Graduate School of Pharmaceutical Sciences, Ewha Womans University, Seoul 03760, Korea

² Department of Pharmaceutical Sciences, University of California, Irvine, CA 92697, USA

nearly white color (Kimura et al. 2008). The characteristic botanical features of *A. keiskei* are described in Flora of Japan:

Rhizomes thick and short, with few elongate roots. *Stems* stout, branched above, 80–120 cm long, glabrous. *Leaves* radical and lower cauline, usually evergreen, 2–3-ternate-pinnate, obtusely deltoid in outline, 20–60 cm. long, glabrous; leaflets rather thick, usually deeply 2–3 cleft of parted; segments usually ovate, 5–10 cm long, 3–6 cm wide, acute, toothed, with impressed veinlets on upper side; sheaths on the upper part elliptic, inflated. *Inflorescence* umbels rather numerous; peduncles in upper part; the inner side of rays and pedicels puberulent; rays 10–20; pedicels 20–40; bracteoles of involucre several, linear or slightly broaden at base, caudately elongate, slightly longer or shorter than pedicels; styles short, as long as stylopodium. *Fruit* oblong, glabrous, 6–8 mm long, flat; dorsal ribs short and rather thick; lateral ribs winged; vittae 1 in intervals, 4 on the commissure (Oi et al. 1965).

Phytochemical investigations on *A. keiskei*

Previous phytochemical investigations on *A. keiskei* have led to the identification of over 100 compounds including various types of flavonoids (1–31, 41–56), coumarins (57–92), phenolic (96), acetylenes (98–102), sesquiterpene (103), diterpene (104), and triterpenes (105–106). Structures of the reported constituents listed in Tables 1, 2 and 3 are shown in Fig. 1. In our recent study on the aerial parts of *A. keiskei*, eight constituents, 4,2',4'-trihydroxy-3'-[(2*E*,5*E*)-7-methoxy-3,7-dimethyl-2,5-octadienyl]chalcone (32), (±)-4,2',4'-trihydroxy-3'-[(2*E*)-6-hydroxy-7-methoxy-3,7-dimethyl-2-octenyl]chalcone (33), 4,2',4'-trihydroxy-3'-[(2*E*)-3-methyl-5-(1,3-dioxolan-2-yl)-2-pentenyl]chalcone (34), 2',3'-furano-4-hydroxy-4'-methoxychalcone (35), (±)-4-hydroxy-2',3'-(2,3-dihydro-2-methoxyfurano)-4'-methoxychalcone (36), 3'-carboxymethyl-4,2'-dihydroxy-4'-methoxychalcone (37), (±)-4,2',4'-trihydroxy-3'-[(3-hydroxy-2,2-dimethyl-6-methylenecyclohexyl)methyl]chalcone (38), and 2''-hydroxyangelichalcone (39) were isolated and reported as new compounds from an ethyl acetate extract along with eight known compounds, xanthoangelol (1), xanthoangelol F (2), 4-hydroxyderricin (3), xanthoangelol G (6), xanthoangelol D (9), xanthoangelol E (10), xanthoangelol H (12), (±)-4,2',4'-trihydroxy-3'-[(6*E*)-2-hydroxy-7-methyl-3-methylene-6-octenyl]chalcone (27), artocarmitin A (40), (+)-*cis*-(3'*R*,4'*R*)-methylkhellactone (93), (-)-*trans*-(3'*R*,4'*S*)-methylkhellactone (94), 3,4-dihydroxanthotoxin (95), and (*Z*)-*p*-coumaryl alcohol (97) (Kil

et al. 2015a, 2016). In particular, xanthoangelol (1) and 4-hydroderricin (3) were isolated in bulk (1: 5 g, 2: 1.8 g), which supported that the two chalcones are the predominant constituents in this herb. As a part of our research on the aerial parts of *A. keiskei*, an effective method was developed to prepare two chalcones from the crude extract using high-speed counter-current chromatography (two-phase solvent system: *n*-hexane–ethyl acetate–methanol–water 9:5:9:4, revolution rate: 800 rpm, flow rate: 1.5 mL/min) (Kil et al. 2015b).

Biological activities of extracts

Anti-inflammatory activity

An *n*-hexane fraction from an ethanol extract of the aerial parts of *A. keiskei* showed inhibitory effects on production of lipopolysaccharide (LPS)-induced nitric oxide (NO) and prostaglandin E₂ (PGE₂) by downregulating inducible NO synthase (iNOS) and cyclooxygenase-2 (COX-2) protein and mRNA (Lee et al. 2010). An ethyl acetate fraction also inhibited production of NO in LPS-activated RAW 264.7 cells (Chang et al. 2014).

Anti-obesity activity

In a study in mice, an ethyl acetate extract was reported to prevent high-fat diet induced adiposity by modulating lipid metabolism through increasing phosphorylation of adenosine monophosphate-activated protein kinase (AMPK) in adipose tissue and liver (Zhang et al. 2015). A patent has been filed on the anti-obesity activity of the ethanol extract of the aerial parts of *A. keiskei* in mice (Lee et al. 2012a).

Anti-hyperlipidemic activity

Dietary *A. keiskei* as an ethyl acetate extract from yellow exudates produced elevation of serum high-density lipoprotein (HDL) levels containing apolipoprotein A1 (ApoA1) and ApoE and reduction of liver triglyceride (TG) levels with decreased mRNA expression of hepatic Acyl-coenzyme A (CoA) synthetase in stroke-prone spontaneously hypertensive rats (SHRSP) (Ogawa et al. 2003). An ethyl acetate fraction from a methanol extract showed inhibitory effects against 3-hydroxy-3-methylglutaryl CoA (HMG-CoA) reductase, a crucial and rate limiting enzyme in biosynthesis of cholesterol (Park et al. 1997b). The impact of *A. keiskei* was examined in rats fed a high fat diet, and it was shown that liver weight was decreased and gene expression of the anti-oxidant enzymes, catalase and glutathione-*s*-reductase, in the liver was increased (Kim et al. 2012). In addition, treatment of the ethanol extract

Table 1 The reported biological activities of xanthoangelol (1)

Part ^a	Biological activity	Reference(s)
RT	First isolation from the part	Kozawa et al. (1977)
RT	Anti-ulcer activity: gastric H ⁺ –K ⁺ ATPase	Murakami et al. (1990)
RT	Anti-bacterial activity: gram-positive bacteria	Inamori et al. (1991)
Syn	Anti-bacterial activity: gram-positive bacteria	Sugamoto et al. (2011)
RT	Anti-tumor-promoter activity: TPA-stimulated ³² Pi-incorporation, Ca ⁺ -calmodulin system	Okuyama et al. (1991)
EXD	Anti-tumor-promoter activity: EBV-EA by TPA	Akihisa et al. (2003)
RT	Anti-tumor and anti-metastatic activities: DNA synthesis, tumor-induced neovascularization, binding of VEGF to HUVECs	Kimura and Baba (2003)
EXD	Anti-tumor activity: apoptosis (caspase-3) in human neuroblastoma (IMR-32), leukemia (Jurkat) cells	Tabata et al. (2005)
EXD	Anti-tumor activity: apoptosis (caspase-9, cytochrome c), ROS, DJ-1 protein in human neuroblastoma (IMR-32) cells	Motani et al. (2008)
EXD, ST	Cytotoxicities: human neuroblastoma cells (LA-N-1, NB-39: drug-resistant; IMR-32, SK-N-SH: drug-sensitive)	Motani et al. (2008), Nishimura et al. (2007)
ST	Anti-tumor activity: apoptosis (DNA fragmentation) in human stomach cancer (KATO III) cells	Takaoka et al. (2008)
AP	Anti-diabetic activity: insulin-like activity	Enoki et al. (2007)
RT	Anti-diabetic activity: GLUT4 translocation	Kawabata et al. (2011)
AP	Anti-diabetic activity: α-glucosidase	Luo et al. (2012)
ST	Anti-diabetic activity: PTP1B	Li et al. (2015)
RT	Anti-hypertensive activity: vasoconstriction in rat aortic rings	Matsuura et al. (2001)
RT	Preventive effect against metabolic syndrome: adiponectin production	Ohnogi et al. (2012b)
RT	Preventive effect against metabolic syndrome: adipocytes differentiation (AMPK, MAPK pathways; C/EBPs, PPARγ)	Zhang et al. (2013)
EXD	Anti-inflammatory activity (details in the patent)	Akihisa et al. (2007)
RT	Anti-inflammatory activity: NO, TNF-α, iNOS, COX-2, AP-1	Yasuda et al. (2014)
LF	Anti-inflammatory activity: IL-6 in TNF-α-stimulated osteosarcoma cells	Shin et al. (2011)
AP	Anti-oxidative activity: NQO1 induction, DPPH radical scavenging	Luo et al. (2012)
ST	Anti-oxidative activity: XO	Kim et al. (2014)
ST	Anti-thrombotic activity: PAI-1	Ohkura et al. (2011)
RT	Anti-thrombotic activity: platelet aggregation (induced by collagen, platelet-activating factor, phorbol 12-myristate 13-acetate, not thrombin)	Son et al. (2014)
ST	Whitening activity: melanin formation in B16 melanoma cells	Arung et al. (2012)
AP	Anti-depressant activity: MAO (nonselective), DBH	Kim et al. (2013)
WP	Anti-viral activity: SARS-CoV (viral proteases: 3CL ^{pro} , PL ^{pro})	Park et al. (2015)

^a Plant part code: *WP* whole plant, *RT* root, *AP* aerial part (including stem and leaf), *ST* stem, *LF* leaf, *EXD* exudate; other source code: *Syn* synthetic

reduced increased levels of serum insulin and triglycerides in rats after drinking fructose. The ethanol extract enhanced expression of genes related to fatty acid β-oxidation and HDL production, acyl-CoA oxidase 1 (ACO1), medium-chain acyl-CoA dehydrogenase (MCAD), ATP-binding membrane cassette transporter A1 (ABCA1), and ApoA1 genes in the fructose fed rats (Ohnogi et al. 2012a).

Anti-oxidative activity

The alcohol extracts of *A. keiskei* were evaluated for their scavenging capability against superoxide anion and 1,1-diphenyl-2-picrylhydrazyl (DPPH) free radicals, and

resulted in decreases of free radical formations. The scavenging effects of the leaves extract were greater than those of the stems extract (Qin et al. 2014; Guo et al. 2013).

Anti-thrombotic activity

Exudates of *A. keiskei* resulted in anti-coagulation by inhibiting elevated plasma plasminogen activator inhibitor 1 (PAI-1) under inflammatory conditions induced by LPS (Ohkura et al. 2011). A methanol extract of its roots showed anti-platelet activity when evaluated in a washed rabbit platelets model system (Son et al. 2014).

Table 2 The reported biological activities of 4-hydroxyderricin (**3**)

Part ^a	Biological activity	Reference(s)
RT	First isolation from the part	Kozawa et al. (1977)
RT	Anti-ulcer activity: gastric H ⁺ -K ⁺ ATPase	Murakami et al. (1990)
RT	Anti-bacterial activity: gram-positive bacteria	Inamori et al. (1991)
Syn	Anti-bacterial activity: gram-positive bacteria	Sugamoto et al. (2011)
Syn	Anti-bacterial activity: <i>S. aureus</i> seryl-tRNA synthetase	Battenberg et al. (2013)
RT	Anti-tumor-promoter activity: TPA-stimulated ³² Pi-incorporation, Ca ⁺ -calmodulin system	Okuyama et al. (1991)
EXD	Anti-tumor-promoter activity: EBV-EA by TPA	Akihisa et al. (2003)
ST	Anti-tumor activity: apoptosis (DNA fragmentation) in human stomach cancer (KATO III) cells	Takaoka et al. (2008)
RT	Anti-tumor and anti-metastatic activities: DNA synthesis, tumor-induced neovascularization, lymphocytes, CD4 ⁺ , CD8 ⁺ , NK-T cells	Kimura et al. (2004)
RT	Anti-tumor activity: apoptosis (DNA topoisomerase, procaspases-3, -8, -9 in human leukemia (HL60) cells Cytotoxicities: human leukemia (HL60), melanoma (CRL1579), lung cancer (A549), stomach cancer (AZ521) cells	Akihisa et al. (2011)
ST	Cytotoxicities: human neuroblastoma cells (IMR-32, NB-39)	Nishimura et al. (2007)
RT	Anti-hypertensive activity: vasoconstriction in rat aortic rings	Matsuura et al. (2001)
ST	Anti-hypertensive activity: systolic blood pressure in SHRSP Anti-hyperlipidemic activity: serum VLDL (microsomal TG transfer-protein), hepatic TG (differentiation factor 1, fatty acid synthase) levels in SHRSP	Ogawa et al. (2005b)
AP	Anti-diabetic activity: insulin-like activity	Enoki et al. (2007)
RT	Anti-diabetic activity: chronic ingestion on borderline mild hyperglycemia (adiponectin)	Ohnogi et al. (2007)
RT	Anti-diabetic activity: GLUT4 translocation	Kawabata et al. (2011)
AP	Anti-diabetic activity: α -glucosidase	Luo et al. (2012)
ST	Anti-diabetic activity: PTP1B	Li et al. (2015)
EXD	Anti-inflammatory activity (details in the patent)	Akihisa et al. (2007)
LF	Anti-inflammatory activity: IL-6 in TNF- α -stimulated osteosarcoma cells	Shin et al. (2011)
RT, AP	Anti-inflammatory activity: NO, TNF- α , iNOS, COX-2, AP-1	Yasuda et al. (2014), Chang et al. (2014)
AP	Anti-inflammatory activity: I- κ B degradation, NF- κ B nuclear translocation	Chang et al. (2014)
RT	Anti-thrombotic activity: platelet aggregation (induced by collagen, platelet-activating factor, phorbol 12-myristate 13-acetate, not thrombin)	Son et al. (2014)
LF	Anti-viral activity: influenza virus neuraminidase	Park et al. (2011)
WP	Anti-viral activity: SARS-CoV (viral proteases: 3CL ^{pro} , PL ^{pro})	Park et al. (2015)
RT	Preventive effect against metabolic syndrome: adiponectin production	Ohnogi et al. (2012b)
RT	Preventive effect against metabolic syndrome: adipocytes differentiation (AMPK, MAPK pathways; C/EBPs, PPAR γ)	Zhang et al. (2013)
AP	Anti-oxidative activity: NQO1 induction	Luo et al. (2012)
WP	Anti-oxidative activity: XO	Kim et al. (2014)
ST	Whitening activity: melanin formation in B16 melanoma cells	Arung et al. (2012)
AP	Anti-depressant activity: inhibition of MAO-B (selective), DBH	Kim et al. (2013)

^a Plant part code: WP whole plant, RT root, AP aerial part (including stem and leaf), ST stem, LF leaf, EXD exudate; other source code: Syn synthetic

Anti-tumor and anti-mutagenic activity

The extracts induced apoptotic cell death in various human cancer cell lines including breast cancer (MDA-MB-231) (Jeong and Kang 2011), leukemia (HL60), melanoma (CRL1579), lung cancer (A549), stomach cancer (AZ521) (Akihisa et al. 2011), prostate cancer (LNCaP) (Lee et al. 2014). In these cancer cell lines

studies, increasing expression level of the proapoptotic gene, Caspase-3, and apoptotic inducer gene, B cell lymphoma 2 (Bcl-2), and decreasing expression level of the apoptotic suppressor gene, Bcl-2 associated x (Bax), were presented as the mechanism of action of anti-cancer effects of *A. keiskei* (Jeong and Kang 2011). The methanol extract showed an anti-mutagenic activity against aflatoxin B1 (AFB1), *N*-methyl-*N'*-nitro-*N*-nitrosoguanidine,

Table 3 Constituents from *A. keiskei* and their reported biological activities

Compound class/name	Part ^a	Biological activity	Reference(s)
<i>Flavonoids</i>			
<i>Chalcones</i>			
Xanthoangelol F (2)	RT	First isolation from the part	Nakata et al. (1999)
	RT	Anti-hypertensive activity: phenylephrine-induced vasoconstriction in rat aortic rings	Matsuura et al. (2001)
	EXD	Anti-tumor-promoter activity: EBV-EA by TPA Anti-tumor-initiator activity: NOR 1	Akihisa et al. (2003)
	EXD	Anti-inflammatory activity (details in the patent)	Akihisa et al. (2007)
	ST	Cytotoxicities: human neuroblastoma (IMR-32, NB-39)	Nishimura et al. (2007)
	Syn	Anti-bacterial activity: Gram-positive bacteria	Sugamoto et al. (2011)
	AP	Anti-viral activity: influenza virus neuraminidase	Park et al. (2011)
	WP	Anti-viral activity: SARS-CoV (3CL ^{pro} , PL ^{pro})	Park et al. (2015)
	WP	Anti-oxidative activity: XO	Kim et al. (2014)
	ST	Anti-diabetic activity: PTP1B	Li et al. (2015)
	RT	First isolation from the part	Nakata et al. (1999)
	EXD	Anti-tumor-promoter activity: EBV-EA by TPA, <i>in vivo</i> Two-stage mouse skin carcinogenesis test Anti-tumor-initiator activity: NOR 1	Akihisa et al. (2003) Akihisa et al. (2006)
	ST	Anti-tumor activity: apoptosis (procaspase-3, -9, Bax) in human neuroblastoma cells (IMR-32, NB-39)	Nishimura et al. (2007)
	EXD	Anti-inflammatory activity (details in the patent)	Akihisa et al. (2007)
	Com	Anti-inflammatory activity: iNOS	Shin et al. (2013)
Isobavachalcone (4)	Syn	Anti-bacterial activity: Gram-positive bacteria	Sugamoto et al. (2011)
	AP	Anti-oxidative activity: NQO1 induction, DPPH radical scavenging	Luo et al. (2012)
	WP	Anti-oxidative activity: XO	Kim et al. (2014)
	AP	Anti-diabetic activity: α -glucosidase	Luo et al. (2012)
	WP	Anti-viral activity: SARS-CoV (3CL ^{pro} , PL ^{pro})	Park et al. (2015)
	RT	First isolation from the part	Baba et al. (1990b)
	RT	Anti-hypertensive activity: phenylephrine-induced vasoconstriction in rat aortic rings	Matsuura et al. (2001)
	RT	Anti-allergic activity: compound 48/80-induced histamine release	Nakata and Baba (2001)
	ST	Anti-oxidative activity: superoxide-scavenging	Aoki et al. (2008)
	WP	Anti-oxidative activity: XO	Kim et al. (2014)
Xanthoangelol B (5)	LF	Anti-inflammatory activity: IL-6 in TNF- α -stimulated osteosarcoma cells	Shin et al. (2011)
	AP	Anti-inflammatory activity: NO, iNOS, COX-2 (I κ B degradation, NF- κ B nuclear translocation)	Chang et al. (2014)
	AP	Anti-viral activity: influenza virus neuraminidase	Park et al. (2011)
	ST	Anti-thrombotic activity: TNF α -induced PAI-1	Ohkura et al. (2011)
	WP	Anti-viral activity: SARS-CoV (3CL ^{pro} , PL ^{pro})	Park et al. (2015)
	RT	First isolation from the part	Nakata et al. (1999)
	AP	Anti-viral activity: influenza virus neuraminidase	Park et al. (2011)
	WP	Anti-viral activity: SARS-CoV (3CL ^{pro} , PL ^{pro})	Park et al. (2015)
	RT	First isolation from the part	Nakata et al. (1999)
	AP	Anti-viral activity: influenza virus neuraminidase	Park et al. (2011)
Xanthoangelol G (6)	WP	Anti-viral activity: SARS-CoV (3CL ^{pro} , PL ^{pro})	Park et al. (2015)
	RT	Preventive effect against metabolic syndrome: adiponectin production	Ohnogi et al. (2012b)
4,2'-Dihydroxy-3'-[(2E)-6-hydroperoxy-3-methyl-7-methylene-2-octenyl]-4'-methoxy chalcone (7)			
	RT	Preventive effect against metabolic syndrome: adiponectin production	Ohnogi et al. (2012b)

Table 3 continued

Compound class/name	Part ^a	Biological activity	Reference(s)
Xanthoangelol C (8)	RT		Baba et al. (1990b)
	RT	Anti-allergic activity: compound 48/80-induced histamine	Nakata and Baba (2001)
Xanthoangelol D (9)	RT	First isolation from the part	Baba et al. (1990b)
	RT	Anti-inflammatory activity: TNF- α induced ET-1 (NF- κ B)	Sugii et al. (2005)
	AP	Anti-viral activity: influenza virus neuraminidase	Park et al. (2011)
	ST	Anti-thrombotic activity: TNF α -induced PAI-1	Ohkura et al. (2011)
	WP	Anti-viral activity: SARS-CoV (3CL ^{pro} , PL ^{pro})	Park et al. (2015)
	ST	Anti-diabetic activity: PTP1B	Li et al. (2015)
Xanthoangelol E (10)	RT	First isolation from the part	Baba et al. (1990b)
	RT	Anti-ulcer activity (details in the patent)	Murakami et al. (1992)
	RT	Anti-thrombotic activity: arachidonic acid metabolism in platelets	Fujita et al. (1992)
	RT	Anti-hypertensive activity: phenylephrine-induced vasoconstriction in rat aortic rings	Matsuura et al. (2001)
	RT	Anti-allergic activity: compound 48/80-induced histamine	Nakata and Baba (2001)
	LF	First isolation from the part	Shin et al. (2011)
	AP	Anti-inflammatory activity: NO, iNOS, COX-2 (I κ B degradation, NF- κ B nuclear translocation)	Chang et al. (2014)
	WP	Anti-viral activity: SARS-CoV (3CL ^{pro} , PL ^{pro})	Park et al. (2015)
	ST	Anti-diabetic activity: PTP1B	Li et al. (2015)
	RT	Anti-tumor-promoter activity: TPA-stimulated ³² Pi-incorporation	Okuyama et al. (1991)
Ashitaba-chalcone (11)	RT	First isolation from the part	Nakata et al. (1999)
	EXD	Anti-tumor-promoter activity: EBV-EA by TPA	Akihisa et al. (2003)
Xanthoangelol H (12)	ST	Cytotoxicities: human neuroblastoma (IMR-32, NB-39)	Nishimura et al. (2007)
	ST	Whitening activity: melanin formation in melanoma cells	Arung et al. (2012)
	EXD	First isolation from the part	Akihisa et al. (2006)
Deoxyxanthoangelol H (13)	EXD	Anti-inflammatory activity (details in the patent)	Akihisa et al. (2007)
	ST	Whitening activity: melanin formation in melanoma cells	Arung et al. (2012)
	EXD	Anti-tumor-promoter activity: EBV-EA by TPA	Akihisa et al. (2006)
Xanthoangelol I (14)	EXD	Anti-tumor-initiator activity: NOR 1	
	ST	Cytotoxicities: human neuroblastoma (IMR-32, NB-39)	Nishimura et al. (2007)
2''-Hydroxy-xanthoangelol I (15)	RT	Protective effect on nerve cells (details in the patent)	Onogi et al. (2004)
	RT	Preventive effect against metabolic syndrome: adiponectin	Ohnogi et al. (2012b)
Xanthoangelol J (16)	EXD	Anti-tumor-promoter activity: EBV-EA by TPA Anti-tumor-initiator activity: NOR 1	Akihisa et al. (2006)
Xanthoangelol K (17)	ST	Anti-diabetic activity: PTP1B	Li et al. (2015)
Xanthoangelol L (18)	ST	First isolation from the part	Li et al. (2015)
Jejuchalcone D (19)	ST	First isolation from the part	Li et al. (2015)
Xanthoangelol M (20)	ST	Anti-diabetic activity: PTP1B	Li et al. (2015)
4-Hydroxy-2',3'-(2,3-dihydro-2-hydroxy isopropylfurano)-4'-methoxychalcone (21)	RT	Preventive effect against metabolic syndrome: adiponectin production	Ohnogi et al. (2012b)
Jejuchalcone E (22)	ST	First isolation from the part	Li et al. (2015)
Dorsmannin A (23)	EXD	First isolation from the part	Akihisa et al. (2006)

Table 3 continued

Compound class/name	Part ^a	Biological activity	Reference(s)
Xanthokeismin A (24)	ST	Anti-oxidative activity: superoxide-scavenging	Aoki et al. (2008)
	AP	Anti-inflammatory activity: NO, iNOS, COX-2 (IκB degradation, NF-κB nuclear translocation)	Chang et al. (2014)
Xanthokeismin B (25)	ST	Anti-oxidative activity: superoxide-scavenging	Aoki et al. (2008)
Xanthokeismin C (26)	ST	Anti-oxidative activity: superoxide-scavenging	Aoki et al. (2008)
4,2',4'-Trihydroxy-3'-[(6 <i>E</i>)-2-hydroxy-7-methyl-3-methylene-6-octenyl]chalcone (27)	LF	Anti-inflammatory activity: IL-6 in TNF-α-stimulated osteosarcoma cells	Shin et al. (2011)
	AP	Protective activity on damaged normal cells: HSPs	Kil et al. (2015a)
Xanthokeistal A (28)	AP	Anti-viral activity: influenza virus neuraminidase	Park et al. (2011)
	WP	Anti-viral activity: SARS-CoV (3CL ^{pro} , PL ^{pro})	Park et al. (2015)
4,2',4'-Trihydroxy-3'-[(2 <i>E</i>)-6,7-dihydroxy-3,7-dimethyl-2-octenyl] chalcone (29)	RT	First isolation from the part	Ohnogi et al. (2012b)
	AP	First isolation from the part	Luo et al. (2012)
4,2'-Dihydroxy-3',4'-[2,3-dihydro-2-[(4 <i>E</i>)-1-hydroxy-1,5-dimethyl-4-hexenyl]furano] chalcone (30)	RT	Preventive effect against metabolic syndrome: adiponectin production	Ohnogi et al. (2012b)
Angelichalcone (31)	RT	Protective effect on nerve cells (details in the patent)	Onogi et al. (2004)
	RT	Osteogenesis promotive activity (details in the patent)	Ohnogi et al. (2004)
4,2',4'-Trihydroxy-3'-[(2 <i>E</i> ,5 <i>E</i>)-7-methoxy-3,7-dimethyl-2,5-octadienyl] chalcone (32)	AP	First isolation from the part	Kil et al. (2015a)
(±)-4,2',4'-Trihydroxy-3'-[(2 <i>E</i>)-6-hydroxy-7-methoxy-3,7-dimethyl-2-octenyl]chalcone (33)	AP	First isolation from the part	Kil et al. (2015a)
4,2',4'-Trihydroxy-3'-[(2 <i>E</i>)-3-methyl-5-(1,3-dioxolan-2-yl)-2-pentenyl]chalcone (34)	AP	First isolation from the part	Kil et al. (2015a)
2',3'-Furano-4-hydroxy-4'-methoxychalcone (35)	AP	First isolation from the part	Kil et al. (2015a)
(±)-4-Hydroxy-2',3'-(2,3-dihydro-2-methoxyfurano)-4'-methoxychalcone (36)	AP	First isolation from the part	Kil et al. (2015a)
3'-Carboxymethyl- 4,2'-dihydroxy-4'-methoxy chalcone (37)	AP	First isolation from the part	Kil et al. (2016)
(±)-4,2',4'-Trihydroxy-3'-[(3-hydroxy-2,2-dimethyl-6-methylenecyclohexyl)methyl]chalcone (38)	AP	First isolation from the part	Kil et al. (2016)
2''-Hydroxy angelichalcone (39)	AP	First isolation from the part	Kil et al. (2016)
Artocarmitin A (40)	AP	First isolation from the part	Kil et al. (2016)
Dihydrochalcones			
Deoxydihydroxanthoan	EXD	First isolation from the part	Akihisa et al. (2006)
Gelol H (41)	ST	whitening activity: melanin formation in melanoma cells	Arung et al. (2012)
Erioschalcone A (42)	AP	First isolation from the part	Luo et al. (2012)
Flavones			
Cynaroside (43) (=luteolin-7- <i>O</i> -β-D-glucopyranoside)	AP	Anti-hyperlipidemic activity	Park et al. (1995)
	AP	Anti-cholesterogenic activity: HMG-CoA reductase	Park et al. (1997b)
	AP	Anti-mutagenic activity (AFB1)	Park et al. (1997a)
	AP	Anti-oxidative activity: epoxide hydrolase in liver	Park et al. (2002)
	AP	Anti-oxidative activity: DPPH radical scavenging	Kim et al. (2005)
	AP	Anti-depressant activity: DBH	Kim et al. (2013)
	ST	Anti-diabetic activity: PTP1B	Li et al. (2015)
Luteolin-7- <i>O</i> -β-D-rutinoside (44)	AP	First isolation from the part	Park et al. (1995)
	AP	Anti-mutagenic activity (AFB1)	Park et al. (1997a)
	AP	Anti-oxidative activity: DPPH radical scavenging	Kim et al. (2005)

Table 3 continued

Compound class/name	Part ^a	Biological activity	Reference(s)
Luteolin (45)	LF	First isolation from the part	Sugamoto et al. (2008)
Flavonol glycosides			
Kaempferol-3- <i>O</i> - α -D-arabinopyranoside (46)	AP	Anti-oxidative activity: DPPH radical scavenging	Kim et al. (2005)
Isoquercitrin (=quercetin-3- <i>O</i> - β -D-glucopyranoside) (47)	AP	Anti-oxidative activity: DPPH radical scavenging	Kim et al. (2005)
Hyperoside (=quercetin-3- <i>O</i> - β -D-galactopyranoside) (48)	AP	First isolation from the part	Park et al. (1996)
	AP	Anti-cholesterogenic activity: HMG-CoA reductase	Park et al. (1997b)
	AP	Anti-mutagenic activity (AFB1)	Park et al. (1997a)
	AP	Anti-oxidative activity: DPPH radical scavenging	Kim et al. (2005)
Quercitrin (49) (=quercetin-3- <i>O</i> - β -D-rhamnopyranoside)	LF	First isolation from the part	Sugamoto et al. (2008)
Quercetin-3- <i>O</i> - α -D-arabinopyranoside (50)	AP	Anti-oxidative activity: DPPH radical scavenging	Kim et al. (2005)
Flavanones			
Munduleaflavanone A (51)	EXD	Anti-tumor-promoter activity: EBV-EA by TPA	Akihisa et al. (2003)
	AP	Anti-diabetic activity: α -glucosidase	Luo et al. (2012)
Munduleaflavanone B (52)	EXD	Anti-tumor-promoter activity: EBV-EA by TPA	Akihisa et al. (2006)
		Anti-tumor-initiator activity: NOR 1	
	EXD	anti-inflammatory activity: TPA-induced mouse ear edema	Akihisa et al. (2007)
Isobavachin (53)	EXD	Anti-tumor-promoter activity: EBV-EA by TPA	Akihisa et al. (2006)
Prostratol F (54)	EXD	Anti-tumor-promoter activity: EBV-EA by TPA	Akihisa et al. (2003)
Sophoraflavanone A (55) (=8-geranylnaringenin)	EXD	Anti-tumor-promoter activity: EBV-EA by TPA	Akihisa et al. (2006)
		Anti-tumor-initiator activity: NOR 1	
	EXD	Anti-inflammatory activity: TPA-induced mouse ear edema	Akihisa et al. (2007)
4'- <i>O</i> -Geranylnaringenin (56)	EXD	Anti-tumor-promoter activity: EBV-EA by TPA	Akihisa et al. (2003)
Coumarins			
Umbelliferone (57)	RT	First isolation from the part	Baba et al. (1990b)
Scopoletin (58)	RT	First isolation from the part	Baba et al. (1990b)
Demethylsuberosin (59)	AP	Anti-diabetic activity: α -glucosidase	Luo et al. (2012)
Osthenol (60)	EXD	Anti-tumor-promoter activity: EBV-EA by TPA	Akihisa et al. (2006)
		Anti-tumor-initiator activity: NOR 1	
	EXD	Anti-inflammatory activity (details in the patent)	Akihisa et al. (2007)
7- <i>O</i> - β -D-Glucopyranosyloxy-8-prenyl coumarin (61)	RT	NGF production enhance effect on in mouse fibroblasts (details in the patent)	Ohnogi et al. (2002)
Psoralen (62)	RT	First isolation from the part	Hata and Kozawa (1961)
Ergapten (63)	AP	First isolation from the part	Luo et al. (2012)
	RT	First isolation from the part	Hata and Kozawa (1961)
Xanthotoxin (64)	FR	First isolation from the part	Baba et al. (1990a)
	AP	First isolation from the part	Luo et al. (2012)
	ST	Anti-diabetic activity: PTP1B	Li et al. (2015)
	RT	First isolation from the part	Hata and Kozawa (1961)
Isopimpinellin (65)	FR	First isolation from the part	Baba et al. (1990a)
	EXD	First isolation from the part	Akihisa et al. (2006)
	AP	First isolation from the part	Luo et al. (2012)
	RT	First isolation from the part	Baba et al. (1990b)
	FR	First isolation from the part	Nakata et al. (1997)
	EXD	First isolation from the part	Akihisa et al. (2006)

Table 3 continued

Compound class/name	Part ^a	Biological activity	Reference(s)
Imperatorin (66)	FR	First isolation from the part	Baba et al. (1990a)
	RT	First isolation from the part	Nakata et al. (1997)
	AP	First isolation from the part	Luo et al. (2012)
Isogosferol (67)	FR	First isolation from the part	Baba et al. (1990a)
Isoimperatorin (68)	FR	First isolation from the part	Baba et al. (1990a)
	RT	First isolation from the part	Nakata et al. (1997)
Saxaline (69)	FR	First isolation from the part	Nakata et al. (1997)
Marmesin (70)	RT	First isolation from the part	Baba et al. (1990b)
(-)-Oxypeucedanin (71)	FR	First isolation from the part	Baba et al. (1990a)
	RT	First isolation from the part	Nakata et al. (1997)
(+)-Oxypeucedanin hydrate (72)	FR,	First isolation from the part	Nakata et al. (1997)
	RT	First isolation from the part	
(-)-Oxypeucedanin hydrate (73)	FR	First isolation from the part	Baba et al. (1990a)
(-)- <i>Tert-O</i> -methoxy peucedanin hydrate (74)	FR	First isolation from the part	Baba et al. (1990a)
(-)- <i>Sec-O</i> -acetoxyp eucedanin hydrate(75)	FR	First isolation from the part	Baba et al. (1990a)
Pseudoisopsoralen (76)	AP	First isolation from the part	Luo et al. (2012)
Angelicin (77)	RT	First isolation from the part	Hata and Kozawa (1961)
Columbianadin (78)	RT	First isolation from the part	Kozawa et al. (1978)
	FR	First isolation from the part	Baba et al. 1990a)
(3' <i>R</i>)-Hydroxy columbianadin (79) (= (8 <i>S</i> ,9 <i>R</i>)-8-angeloyl oxy-8,9-dihydrooroselol)	EXD	Anti-tumor-promoter activity: EBV-EA by TPA	Akihisa et al. (2003)
	RT	First isolation from the part	Sugamoto et al. (2009)
(8 <i>S</i> ,9 <i>R</i>)-9-Angeloyloxy-8,9-dihydrooroselol (80)	RT	First isolation from the part	Kozawa et al. (1978)
	RT	Anti-tumor-promoter activity: TPA-stimulated ³² Pi-incorporation	Okuyama et al. (1991)
Archangelicin (81)	RT	First isolation from the part	Kozawa et al. (1978)
	FR	First isolation from the part	Baba et al. (1990a)
	RT	Anti-tumor-promoter activity: TPA-stimulated ³² Pi-incorporation	Okuyama et al. (1991)
Lomatin (82)	FR	First isolation from the part	Baba et al. (1990a)
Selinidin (83)	RT	First isolation from the part	Baba et al. (1990b)
	FR	First isolation from the part	Baba et al. (1990a)
	EXD	Anti-tumor-initiator activity: NOR 1	Akihisa et al. (2003)
	EXD	Anti-allergic activity: IgE-mediated mast cell activation (β-hexosaninidase, LTC ₄ , TNF-α, FcεRI β-chain PLCγ1, p38 MAPK, IkB-α)	Kishiro et al. (2008)
	AP	First isolation from the part	Luo et al. (2012)
	ST	Anti-diabetic activity: PTP1B	Li et al. (2015)
Selidinin (84)	EXD	Anti-inflammatory activity (details in the patent)	Akihisa et al. (2007)
<i>O</i> -Isobutyroyllomatin (85)	ST	Anti-diabetic activity: PTP1B	Li et al. (2015)
Khellactone (86)	AP	First isolation from the part	Luo et al. (2012)
	ST	Anti-diabetic activity: PTP1B	Li et al. (2015)
3'-Seneciroyl khellactone (87)	EXD	Anti-tumor-promoter activity: EBV-EA by TPA	Akihisa et al. (2003)
		Anti-tumor-initiator activity: NOR 1	
	AP	Anti-diabetic activity: α-glucosidase	Luo et al. (2012)

Table 3 continued

Compound class/name	Part ^a	Biological activity	Reference(s)
Peguanguenine (88)	EXD	Anti-inflammatory activity: TPA-induced mouse ear edema	Akihisa et al. (2007)
(3' <i>R</i> ,4' <i>R</i>)-3'-Angeloyl khellactone (89) (miswritten as isolaserpiten)	RT	First isolation from the part	Baba et al. (1990b)
	FR	First isolation from the part	Baba et al. (1990a)
	EXD	Anti-tumor-promoter activity: EBV-EA by TPA Anti-tumor-initiator activity: NOR 1	Akihisa et al. (2003)
	EXD	Anti-inflammatory activity (details in the patent)	Akihisa et al. (2007)
4'-Senecioid khellactone (90)	EXD	Anti-tumor-promoter activity: EBV-EA by TPA Anti-tumor-initiator activity: NOR 1	Akihisa et al. (2003)
Isolemanidin (91) (miswritten as laserpiten)	RT	First isolation from the part	Baba et al. (1990b)
	FR	First isolation from the part	Baba et al. (1990a)
	EXD	Anti-tumor-promoter activity: EBV-EA by TPA Anti-tumor-initiator activity: NOR 1	Akihisa et al. (2003)
	ST	Anti-hyperlipidemic activity: serum HDL↑, hepatic TG↓ (hepatic TG lipase, differentiation factor 1)	Ogawa et al. (2005b)
	EXD	Anti-inflammatory activity: TPA-induced mouse ear edema	Akihisa et al. (2007)
Pteryxin (92)	EXD	Anti-tumor-promoter activity: EBV-EA by TPA Anti-tumor-initiator activity: NOR 1	Akihisa et al. (2003)
	EXD	Anti-inflammatory activity (details in the patent)	Akihisa et al. (2007)
(+)- <i>Cis</i> -(3' <i>R</i> ,4' <i>R</i>)-methylkhellactone (93)	AP	First isolation from the part	Kil et al. (2016)
(-)- <i>Trans</i> -(3' <i>R</i> ,4' <i>S</i>)-methylkhellactone (94), Dihydrocoumarin	AP	First isolation from the part	Kil et al. (2016)
3,4-Dihydroxanthotoxin (95)	AP	First isolation from the part	Kil et al. (2016)
<i>Phenolics</i>			
5- <i>n</i> -Pentadecylresorcinol (96)	EXD	First isolation from the part	Akihisa et al. (2003)
(<i>Z</i>)- <i>p</i> -Coumaryl alcohol (97)	AP	First isolation from the part	Kil et al. (2016)
<i>Acetylenes</i>			
Falcarindiol (98)	EXD	First isolation from the part	Akihisa et al. (2003)
	AP	Anti-diabetic activity: α-glucosidase	Luo et al. (2012)
	ST	Anti-diabetic activity: PTP1B	Li et al. (2015)
<i>Acetylenes</i>			
(11 <i>S</i> ,16 <i>R</i> , <i>Z</i>)-11,16-Dihydroxyoctadeca-9,17-dien-12,14-diynyl acetate (99)	EXD	First isolation from the part	Akihisa et al. (2006)
(10 <i>S</i> ,15 <i>R</i> , <i>Z</i>)-10,15-Dihydroxyheptadeca-8,16-dien-11,13-diynyl acetate (100)	AP	Anti-diabetic activity: α-glucosidase	Luo et al. (2012)
(11 <i>S</i> ,16 <i>R</i> , <i>Z</i>)-Octadeca-9,17-dien-12,14-diyne-1,11,16-triol (101)	AP	First isolation from the part	Luo et al. (2012)
Angelicol B (102)	AP	First isolation from the part	Luo et al. (2012)
<i>Sesquiterpene</i>			
Ashitabaol A (103)	SD	Anti-oxidative: free radical scavenging	Aoki and Ohta (2010)
<i>Diterpene</i>			
Steviol-13- <i>O</i> -β-D-glucopyranoside 19-β-D-glucopyranosyl ester (104)	LF	First isolation from the part	Zhou et al. (2012)
<i>Triterpenes</i>			
Daucosterol (105)	LF	First isolation from the part	Zhou et al. (2012)
Stigmasterol (106)	LF	First isolation from the part	Zhou et al. (2012)
<i>Triterpenes</i>			
Pregnenolone (107)	AP	Anti-oxidative activity: DPPH radical scavenging	Luo et al. (2012)
<i>Nucleoside</i>			

Table 3 continued

Compound class/name	Part ^a	Biological activity	Reference(s)
Adenosine (108)	AP	First isolation from the part	Park et al. (1996)
Saccharide			
Sucrose (109)	AP	First isolation from the part	Park et al. (1996)
	AP	Anti-mutagenic activity (AFB1)	Park et al. (1997a)
Acids			
Angelic acid (110)	RT	First isolation from the part	Hata and Kozawa (1961)
Behenic acid (111)	RT	First isolation from the part	Hata and Kozawa (1961)
Others			
1-Ceritol (112)	LF	First isolation from the part	Zhou et al. (2012)
(<i>E</i>)-2,6-Dimethylocta-5,7-diene-2,3-diol (113)	AP	First isolation from the part	Luo et al. (2012)
Angelicol A (114)	AP	Anti-oxidative activity: DPPH radical scavenging	Luo et al. (2012)

^a Plant part code: *WP* whole plant, *RT* root, *AP* aerial part (including stem and leaf), *ST* stem, *LF* leaf, *EXD* exudate, *FR* fruit, *SD* seed; other source code: *Syn* synthetic, *Com* commercial

and 4-nitroquinoline-1-oxide in a salmonella assay system (Park et al. 1997a).

Anti-diabetic activity

The ethanol extract exhibited insulin-like activities through a pathway independent of peroxisome proliferator-activated receptor- γ (PPAR- γ) activation (Enoki et al. 2007). Long-term ingestion of a powder of *A. keiskei* improved blood glucose control and moderately and safely reduced blood glucose (Ohnogi et al. 2007). A patent is filed on the beneficial effects of *A. keiskei* on diabetes (Lee et al. 2013).

Anti-bacterial activity

An exudate of *A. keiskei* showed anti-bacterial activity against *Helicobacter pylori*, suggesting *A. keiskei* as a preventive agent for gastritis (Fukuo et al. 2005). A patent on this exudate is filed for controlling mouth odor by an anti-bacterial activity against *Fusobacterium nucleatum* (Saeki et al. 2013).

Hepato-protective activity

A clinical trial for habitual alcohol drinkers with abnormal liver function tests also showed that intake of the extracts significantly reduced γ -glutamyl transferase (GGT) levels (Noh et al. 2015). A patent is filed on the beneficial activity of an ethanol extract of the roots to improve liver functions and relieve fatigue (Kim et al. 2006).

Other biological activities

In addition to these activities, an ethanol extract also showed an anti-hypertensive activity by inhibiting angiotensin 1-converting enzyme (ACE) in spontaneously hypertensive rats, similar to nicotianamine (Shimizu et al. 1999). Green juice of *A. keiskei* was proven to protect lymphocytic DNA damage of smokers, which was reported to potentially reduce the risk of lung cancer in a clinical trial (Kang et al. 2004). In another clinical study, consuming a drink powder of the leaves three times a day for 2 weeks alleviated mild constipation and improved defecation (Kusaba et al. 2008). Deficit of learning and memory caused by Alzheimer's disease and aging may be alleviated by *A. keiskei* since it has been shown that the plant is capable of restoring scopolamine-reduced phosphorylation of cyclic adenosine monophosphate (cAMP) response element-binding protein (CREB) and expression of brain-derived neurotrophic factor (BDNF) in the mouse hippocampus (Oh et al. 2013). Topical uses of the extracts were shown to whiten skin with tyrosinase inhibitory effect (Higa 1985; Lee et al. 2012b) and promote hair growth (Sakaguchi et al. 1996) in patent applications.

Biological activities of isolates from *A. keiskei*

Isolates from *A. keiskei* have showed diverse biological activities. Especially two major chalcones, xanthoangelol (**1**) and 4-hydroxyderricin (**3**), have been extensively




Fig. 1 Chemical structures of compounds isolated from *A. keiskei*




Fig. 1 continued

studied and reported to have anti-tumor, anti-inflammatory, anti-depressant, anti-obesity, anti-thrombotic, anti-diabetic, anti-hyperlipidemic, anti-hypertensive, and anti-ulcer effects. The various effects of isolates will be discussed in

the subsequent sections and summarized in Tables 1, 2 and 3. In our recent research, several compounds from *A. keiskei* were evaluated for their effects on heat shock protein (HSP) expression. HSPs are known to have

protective effects on damaged cells, and HSP-inducible agents may act as prospective agents to protect normal cells from chemotherapy induced adverse effects. (\pm)-4,2',4'-Trihydroxy-3'-[(6*E*)-2-hydroxy-7-methyl-3-methylene-6-octenyl] chalcone (**27**) of 2 μ M increased expressions of HSF1 (4.3-fold), HSP70 (1.5-fold), and HSP27 (4.6-fold) proteins at 6 h of treatment and mRNA levels of HSP70 (5.0-fold) and HSP27 (1.5-fold) at 3 h with lower cytotoxicity than celastrol, a positive control (Kil et al. 2015a).

Anti-ulcer activity

Xanthoangelol (**1**) and 4-hydroxyderricin (**3**) inhibited H^+ , K^+ -ATPase of gastric microsomal vesicles which were prepared from pig fundic mucosa. Additionally, they downregulated K^+ -stimulated *p*-nitrophenyl phosphatase. Xanthoangelol (**1**) reduced the formation of stress-induced gastric lesion as well as acid secretion in rats (Murakami et al. 1990). Xanthoangelol E (**10**) was also found to have the anti-ulcer activity as patented by Murakami et al. (Murakami et al. 1992).

Anti-bacterial activity

Xanthoangelol (**1**) and 4-hydroxyderricin (**3**) showed antibacterial activities against gram-positive bacteria, *Bacillus subtilis* PCI-219, *B. subtilis* ATCC-6633, *B. cereus* FDA-5, *Staphylococcus aureus* 209-P, *S. aureus* IFO-3060, *S. aureus* IFO-3762, and *Micrococcus luteus* IFO-12708, but not gram-negative bacteria. Although the antibacterial activities were lower than gentamicin for the aforementioned bacteria, the anti-bacterial activity of xanthoangelol (**1**) against *M. luteus* IFO-12708 was as strong as that of gentamicin (MIC 0.76 μ g/mL) (Inamori et al. 1991). Mechanism of anti-bacterial activity of 4-hydroxyderricin (**3**) against *S. aureus* was elucidated by utilizing a chemical cleavable linker due to its heat sensitivity, reporting that the isolate inhibits amino acylation of tRNAs catalyzed by *S. aureus* seryl-tRNA synthetase which is an essential enzymatic pathway for bacterial viability (Battenberg et al. 2013). The two chalcones exhibited antibacterial activities against plant-pathogenic bacteria, *Agrobacterium tumefaciens* IFO-3058, *Pseudomonas syringae* pv. *phaseolicola* IFO-12656, *P. syringae* pv. *tabaci* IFO-3508, and *P. stutzeri* IFO-12510 (Inamori et al. 1991). Sumanoto et al. reported anti-bacterial activities of xanthoangelol (**1**), xanthoangelol F (**2**), 4-hydroxyderricin (**3**), and isobavachalcone (**4**), against gram-positive bacteria, *Escherichia coli* NBRC 3301, *P. mirabilis* NBRC 13300, *P. fluorescens* NBRC 3757, *B. subtilis* NBRC 3757, *S. epidermidis* NBRC 12993, and *M. luteus* NBRC 3333 (Sugamoto et al. 2011).

Chemopreventive activity

Bioactive compounds with anti-tumor-promoter, anti-tumor-initiator, and anti-mutagenic activities may be considered as chemopreventive therapeutic agents.

Two angular furanocoumarin, (8*S*,9*R*)-8-angeloxy-8,9-dihydrooroseol (**79**, (3'*R*)-hydroxycolumbianadin) and archangelicin (**81**), and three chalcones, xanthoangelol (**1**), 4-hydroxyderricin (**3**), and ashitaba-chalcone (**11**) inhibited 12-*O*-tetradecanoylphorbol-13-acetate (TPA)-stimulated 32 Pi-incorporation of phospholipids of cultured cells. Xanthoangelol (**1**) and 4-hydroxyderricin (**3**) showed anti-tumor-promoting activities in mouse skin carcinogenesis induced by 7,12-dimethylbenz[α]anthracene (DMBA) plus TPA via the modulation of Ca^{2+} -calmodulin system (Okuyama et al. 1991). Isobavachalcone (**4**) exhibited inhibitory effects on skin tumor promotion in in vivo two-stage mouse skin carcinogenesis test (Akihisa et al. 2006). Inhibitory effects on the induction of Epstein-Barr virus early antigen (EBV-EA) by TPA in Raji cells were found for xanthoangelol (**1**), xanthoangelol F (**2**), 4-hydroxyderricin (**3**), isobavachalcone (**4**), xanthoangelol H (**12**), xanthoangelol I (**14**), xanthoangelol J (**16**), munduleaflavanone A (**51**), munduleaflavanone B (**52**), isobavachin (**53**), prostratol F (**54**), 8-geranylnaringenin (**55**, 8-geranylarnaringenin), 4'-*O*-geranylnaringenin (**56**), osthenol (**60**), (3'*R*)-hydroxycolumbianidin (**79**, (8*S*,9*R*)-8-angeloxy-8,9-dihydrooroseol), 3'-senecioidyl khellactone (**87**), (3'*R*,4'*R*)-3'-angeloyl khellactone (**89**, miswritten as "isolaserapitin" in the article), 4'-senecioidyl khellactone (**90**), isolemanidin (**91**, miswritten as "laserapitin" in the article), and pteryxin (**92**) (Akihisa et al. 2003, 2006). Xanthoangelol F (**2**), isobavachalcone (**4**), xanthoangelol I (**14**), xanthoangelol J (**16**), munduleaflavanone B (**52**), 8-geranylnaringenin (**55**, sophoraflavanone A), osthenol (**60**), selinidin (**83**), 3'-senecioidyl khellactone (**87**), (3'*R*,4'*R*)-3'-angeloyl khellactone (**89**, miswritten as "isolaserapitin" in the article), 4'-senecioidyl khellactone (**90**), isolemanidin (**91**, miswritten as "laserapitin" in the article), and pteryxin (**92**) inhibited activation of (\pm)-(*E*)-methyl-2-[(*E*)-hydroxy-imino]-5-nitro-6-methoxy-3-hexemide (NOR 1), a NO donor (Akihisa et al. 2003, 2006). Cynaroside (**43**, luteolin-7-*O*- β -D-glucopyranoside), luteolin-7-*O*- β -D-rutinoside (**44**), hyperoside (**48**, quercetin-3-*O*- β -D-galactopyranoside), and sucrose (**109**) showed anti-mutagenic activities against aflatoxin B1 (AFB1) (Park et al. 1997a).

Anti-tumor activity

In a previous study, xanthoangelol (**1**) and 4-hydroxyderricin (**3**) were identified as active compounds to inhibit tumor growth in Lewis lung carcinoma (LLC)-bearing mice and to prolong survival time and to suppress

metastasis to the lung after surgical removal of primary tumors from an active ethyl acetate fraction of a 50% ethanol extract (Kimura and Baba 2003; Kimura et al. 2004). The anti-tumor activities of xanthoangelol (1) might be derived from its inhibitory effects on DNA synthesis in LLC cells and on tumor-induced neovascularization by reducing formation of capillary-like tubes by vascular endothelial cells and adhesion of vascular endothelial growth factor (VEGF) to human umbilical vein endothelial cells (HUVECs) (Kimura and Baba 2003). 4-Hydroxyderricin (3) also inhibited the DNA synthesis in LLC cells and the formation of capillary-like tubes, but did not reduce adhesion of vascular endothelial growth factor (VEGF) to HUVECs. Instead, it suppressed reduction of lymphocytes, CD4⁺, CD8⁺, and natural killer (NK)-T cells in spleen of tumor-removed mice (Kimura et al. 2004). Xanthoangelol (1) induced apoptotic cell death by activation of caspase-3 in neuroblastoma (IMR-32) and leukemia (Jurkat) cells through a mechanism, independent of Bax/Bcl-2 signal transduction (Tabata et al. 2005). Their ongoing study revealed details of the mechanism of action of xanthoangelol (1) inducing apoptosis by triggering oxidative stress through generating reactive oxygen species and downregulating antioxidant DJ-1 protein in IMR-32 cells (Motani et al. 2008). Xanthoangelol (1) showed cytotoxic effects on several neuroblastoma cell lines including drug-resistant LA-N-1 and NB-39 cells as well as drug-sensitive IMR-32 and SK-N-SH cells (Nishimura et al. 2007; Motani et al. 2008). Xanthoangelol F (2), 4-hydroxyderricin (3), isobavachalcone (4), xanthoangelol H (12), and xanthoangelol I (14) were cytotoxic to the NB-39 and IMR-32 cells. Isobavachalcone (4) was found to induce apoptotic cell death in the neuroblastoma cells via the mitochondrial pathway including reduction of procaspase-3 and procaspase-9 and induction of Bax level with no cytotoxicity against normal cells (Nishimura et al. 2007). Xanthoangelol (1) and 4-hydroxyderricin (3) inhibited growth of human stomach cancer (KATO III) cells through induction of apoptosis by fragmenting DNA (Takaoka et al. 2008). 4-Hydroxyderricin (3) exhibited cytotoxic activities in human leukemia (HL60), melanoma (CRL1579), lung cancer (A549), and stomach cancer (AZ521) cells. The cytotoxicity to HL60 cells was caused by induction of apoptosis through inhibiting DNA topoisomerase and reducing procaspases-3, -8, and -9 (Akihisa et al. 2011).

Anti-diabetic activity

In a genetically diabetic KK-A^y mouse model, xanthoangelol (1) and 4-hydroxyderricin (3) showed insulin-like activities to enhance glucose uptake and induce pre-adipocyte differentiation into adipocytes without activating PPAR- γ . 4-Hydroxyderricin (3) also prevented the

progression of diabetes in these mice (Enoki et al. 2007). On the basis of the preventive effect, a clinical trial was performed with *A. keiskei* powder containing 4-hydroxyderricin (3), resulting in reduction of blood glucose and improvement of blood glucose control through increasing adiponectin in subjects with borderline or mild hyperglycemia after 12 weeks of consumption of *A. keiskei* (Ohnogi et al. 2007). Xanthoangelol (1), 4-hydroxyderricin (3), isobavachalcone (4), munduleaflavanone A (51), demethylsuberosin (59), 3'-senecioidyl khellactone (87), faltarindiol (98), and (10*S*,15*R*,*Z*)-10,15-dihydroxyheptadeca-8,16-dien-11,13-diynyl acetate (100) appeared to inhibit α -glucosidase, which involved in glucose metabolism (Luo et al. 2012). Xanthoangelol (1) and 4-hydroxyderricin (3) stimulated skeletal muscle-associated glucose uptake by inducing glucose transporter 4 (GLUT-4) translocation, resulting in normal blood glucose level (Kawabata et al. 2011). Recently, protein tyrosine phosphatase 1B (PTP1B) has been suggested as a therapeutic target for treatment of type 2 diabetes based on its down-regulating effect on insulin signaling. Li et al. evaluated isolates from the stems of *A. keiskei* for their PTP1B inhibitory effects using oleanolic acid as a positive control. Xanthoangelol (1), xanthoangelol F (2), 4-hydroxyderricin (3), xanthoangelol D (9), xanthoangelol E (10), xanthoangelol K (17), xanthoangelol M (20) and bergapten (63, 5-methoxypsoralen; miswritten as "methoxsalen" in the article) showed strong PTP1B inhibitory effect (IC₅₀ 0.82–4.42 μ g/mL), together with moderately active four compounds (selinidin, 83; *O*-isobutyroylmatin, 85; khellactone, 86; faltarindiol, 98, IC₅₀ 4.42–11.55 μ g/mL) and a weakly active one (cynaroside, 43, IC₅₀ 17.49 μ g/mL). Furthermore, in a kinetic study, xanthoangelol (1) with IC₅₀ values of 0.82 μ g/mL exhibited typical characteristics of the competitive PTP1B inhibitor and in molecular docking simulations, it was suggested that the ring B of the chalcone skeleton may anchor in a pocket of PTP1B (Li et al. 2015).

Nerve cell protective activity

Onogi et al. filed a patent application presenting angelichalcone (31) with its nerve cell protective effect (Onogi et al. 2004). In addition, 7-*O*- β -D-glucopyranosyloxy-8-prenyl coumarin (61) was reported for its enhancing effect on nerve growth factor (NGF) production in mouse fibroblast L-M cells in a patent (Ohnogi et al. 2002).

Anti-hyperlipidemic activity

Cynaroside (43, luteolin-7-*O*- β -D-glucopyranoside) showed hypolipidemic activity when administrated by intraperitoneal injection in rats (Park et al. 1995).

Cynaroside (**43**) and hyperoside (**48**, quercetin-3-*O*- β -D-galactopyranoside) exhibited anti-cholesterogenic activity by inhibiting HMG-CoA reductase (Park et al. 1997b). When SHRSP was fed diets containing 4-hydroxyderricin for 7 weeks, serum very low-density lipoprotein (VLDL) and hepatic triglycerides (TG) levels were significantly reduced, without any effect on serum HDL levels. It was considered that microsomal TG transfer-protein may be responsible for the decrease in serum VLDL levels and differentiation factor 1 and fatty acid synthase may induce the decrease in hepatic TG content (Ogawa et al. 2005b). In case of isolemanidin (**91**, miswritten as “laserapitin” in the article), the serum HDL levels, especially apoE-HDL, increased and the hepatic TG content decreased, which might be relevant with a decrease in hepatic TG lipase and differentiation factor 1, respectively (Ogawa et al. 2005a).

Preventive effects against metabolic syndrome

Ohnogi et al. isolated six new chalcones, 4,2'-dihydroxy-3'-[(2*E*)-6-hydroperoxy-3-methyl-7-methylene-2-octenyl]-4'-methoxychalcone (**7**), 2''-hydroxyxanthoangelol I (**15**), 4-hydroxy-2',3'-(2,3-dihydro-2-hydroxyisopropylfuran)-4'-methoxychalcone (**21**), 4,2',4'-trihydroxy-3'-[(2*E*)-6,7-dihydroxy-3,7-dimethyl-2-octenyl]chalcone (**29**), 4,2'-dihydroxy-3',4'-[2,3-dihydro-2-[(4*E*)-1-hydroxy-1,5-dimethyl-4-hexenyl]furan]chalcone (**30**), and angelichalcone (**31**) together with xanthoangelol (**1**) and 4-hydroxyderricin (**3**), and evaluated their effects on adiponectin mRNA and protein levels. As a result, all isolates, except for angelichalcone (**31**), increased expression of adiponectin mRNA and production of adiponectin in 3T3-L1 adipocytes. The effects on adiponectin could be expected to prevent metabolic syndrome through improving insulin resistance and dyslipidemia (Ohnogi et al. 2012b). Xanthoangelol (**1**) and 4-hydroxyderricin (**3**) suppressed adipocytes differentiation via AMPK and mitogen-activated protein kinase (MAPK) pathways, which resulted in downregulating expression of adipocyte-specific transcription factors, CCAAT/enhancer-binding protein- β (C/EBP- β), C/EBP- α , and PPAR- γ . Inhibitory effects on adipocytes differentiation could provide preventive effects against obesity and obesity-related disorders (Zhang et al. 2013).

Anti-hypertensive activity

Xanthoangelol (**1**), xanthoangelol F (**2**), 4-hydroxyderricin (**3**), xanthoangelol B (**5**), and xanthoangelol E (**10**) inhibited phenylephrine-induced vasoconstriction. Among them, xanthoangelol (**1**), xanthoangelol F (**2**), 4-hydroxyderricin (**3**), and xanthoangelol E (**10**) showed the inhibitory effects through endothelium-dependent production of endothelium-derived relaxing factor (EDRF)/NO. On the other

hand, xanthoangelol B (**5**) directly inhibited smooth muscle functions through reducing intracellular free calcium $[Ca_2^+]_i$ elevation induced by phenylephrine without affecting EDRF/NO production. 4-Hydroxyderricin (**3**) also suppressed elevation of $[Ca_2^+]_i$ induced by phenylephrine (Matsuura et al. 2001). Dietary 4-hydroxyderricin (**3**) reduced elevation of systolic blood pressure in SHRSP, which also supported its anti-hypertensive activity (Ogawa et al. 2005b).

Anti-allergic activity

Xanthoangelol B (**5**), xanthoangelol C (**8**), and xanthoangelol E (**10**) inhibited the compound 48/80-induced histamine release from rat peritoneal mast cells, whereas, xanthoangelol (**1**) and 4-hydroxyderricin (**3**) enhanced it. Compound 48/80 is one of immunological stimulators, which activates the mast cell to induce allergic responses by mediates (Nakata and Baba 2001). Selinidin (**83**) was found to suppress degranulation, leukotriene C4 (LTC4) synthesis, and tumor necrosis factor- α (TNF- α) production without affecting binding of immunoglobulin E (IgE) with high affinity IgE receptor (Fc ϵ RI) and expression of cell surface Fc ϵ RI in antigen-stimulated primary mouse bone marrow-derived mast cells (BMMCs). Selinidin (**83**) inhibited tyrosine phosphorylation of Fc ϵ RI β -chain and phospholipase C γ 1 (PLC γ 1) not Fc ϵ RI γ -chain in BMMCs upon Fc ϵ RI stimulation, demonstrating its inhibitory effects on Fc ϵ RI-dependent signaling pathway. Phosphorylation of p38 MAPK and inhibitory κ B- α (I κ B- α) were also downregulated by selinidin (**83**) (Kishiro et al. 2008).

Anti-inflammatory activity

Xanthoangelol D (**9**) inhibited activation of nuclear factor- κ B (NF- κ B), which was involved in attenuation of basal and TNF- α -induced endothelin-1 (ET-1), a potent vasoconstrictor peptide. Its inhibitory effect against NF- κ B inflammatory signaling pathway could be useful for treating vascular and inflammatory diseases because many vascular diseases are closely related to the inflammatory processes (Sugii et al. 2005). A patent covers the anti-inflammatory activities of seven flavonoids, xanthoangelol (**1**), xanthoangelol F (**2**), 4-hydroxyderricin (**3**), isobavachalcone (**4**), deoxyxanthoangelol H (**13**), munduleaflavanone B (**52**), and sophoraflavanone A (**55**, 8-geranylnaringenin), and six coumarins, osthonol (**60**), selinidin (**84**), peguanguenine (**88**), (3'*R*,4'*R*)-3'-angeloyl khellactone (**89**), isolemandin (**91**), and pteryxin (**92**) from exudates of *A. keiskei* (Akihisa et al. 2007). Shin et al. reported that xanthoangelol (**1**), 4-hydroxyderricin (**3**), xanthoangelol B (**5**), and 4,2',4'-trihydroxy-3'-[(6*E*)-2-hydroxy-7-methyl-3-methylene-6-octenyl]chalcone (**27**) showed inhibitory

activities of IL-6 production in TNF- α -stimulated MG-63 osteosarcoma cells and they have C-4' hydroxyl groups in common (Shin et al. 2011). Toll-like receptors (TLRs) are essential for induction of innate immune responses against invading pathogens, however, overexpression of transcription factors and pro-inflammatory genes by activation of TLR signaling pathway might cause certain inflammatory diseases. Isobavachalcone (4) reduced overexpression of iNOS induced by TLR agonists (lipopeptide 2-kDa, polyribonucleic polyribocytidylic acid, or LPS) in murine macrophages (Shin et al. 2013). Xanthoangelol (1) and 4-hydroxyderricin (3) reduced production of LPS-induced NO and TNF- α , expression of iNOS and COX-2 in RAW264 mouse macrophages. Moreover, both suppressed DNA binding activities of activator protein-1 (AP-1) and NF- κ B and phosphorylation level of p65 subunit of NF- κ B. However, two chalcones downregulated only AP-1 not NF- κ B (Yasuda et al. 2014). Inhibitory effects on LPS-induced NO and expression of iNOS and COX-2 genes were also found for xanthoangelol B (5), xanthoangelol E (10), and xanthokeismin A (24) as well as 4-hydroxyderricin (3) and their mechanisms involved inhibition of I κ B degradation and NF- κ B nuclear translocation (Chang et al. 2014).

Anti-oxidative activity

Cynaroside (43, luteolin-7-O- β -D-glucopyranoside) inhibited bromobenzene-induced hepatic lipid peroxidation by restoring epoxide hydrolase activity in liver of rats (Park et al. 2002). Xanthoangelol B (5), xanthokeismin A (24), xanthokeismin B (25), and xanthokeismin C (26) showed strong superoxide-scavenging activities. Xanthoangelol (1), 4-hydroxyderricin (3), and isobavachalcone (4) induced reduced nicotinamide adenine dinucleotide (phosphate) (NAD(P)H):quinone oxidoreductase 1 (NQO1) which protects against quinone-mediated oxidative damage. In addition, xanthoangelol (1), isobavachalcone (4), cynaroside (43, luteolin-7-O- β -D-glucopyranoside), luteolin-7-O- β -D-rutinoside (44), kaempferol-3-O- α -D-arabinopyranoside (46), isoquercitrin (47, quercetin-3-O- β -D-glucopyranoside), hyperoside (48, quercetin-3-O- β -D-galactopyranoside), quercetin-3-O- α -D-arabinopyranoside (50), pregnenolone (107), and angelicol A (114) showed DPPH radical scavenging activities (Luo et al. 2012; Kim et al. 2005). Ashitabaol A (103), an isolate from the seeds of *A. keiskei*, was found to be more abundant in the seed coat during germination through HPLC analysis. Ashitabaol A (103) exhibited 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid) (ABTS) free radical scavenging activity and was expected to protect the seed against oxidative damage induced by free radicals during germination (Aoki and Ohta 2010). Xanthine oxidase (XO) leads to formation of free radicals, and cause oxidative damage

to living tissues. Xanthoangelol (1), xanthoangelol F (2), 4-hydroxyderricin (3), isobavachalcone (4), and xanthoangelol B (5) showed inhibitory effects on XO activity, which could be relevant with attenuation of inflammation (Kim et al. 2014).

Anti-thrombotic activity

Xanthoangelol E (10) was evaluated for effects on arachidonic acid metabolism in gastric antral mucosa and platelet of rabbit. Xanthoangelol E (10) showed no significant effect on production of PGE₂, PGF₂ α , and their metabolites in gastric antral mucosal slices, whereas, suppressed production of thromboxane B₂ (TXB₂) and 12-hydroxy-5,8,10-heptadecatrienoic acid (HHT) from exogenous arachidonic acid in platelets (Fujita et al. 1992). Ohkura et al. reported that xanthoangelol (1), xanthoangelol B (5), and xanthoangelol D (9) isolated from the exudates of *A. keiskei* suppressed production of TNF α -induced PAI-1, an inhibitor of fibrinolysis, in human umbilical vein endothelial cells (HUVECs) (Ohkura et al. 2011). Regarding to anti-thrombotic activities, xanthoangelol (1) and 4-hydroxyderricin (3) also exhibited inhibitory effects on platelet aggregation induced by collagen, platelet-activating factor, and phorbol 12-myristate 13-acetate not thrombin (Son et al. 2014).

Anti-depressant activity

Xanthoangelol (1) was compared to iproniazid, an anti-depressant nonselective MAO inhibitor. Xanthoangelol (1) inhibited monoamine oxidase (MAO) in a nonselective manner and suppressed dopamine β -hydroxylase (DBH). On the other hands, 4-hydroxyderricin selectively inhibited MAO-B and mildly suppressed DBH, which was higher than deprenyl, a selective MAO-B inhibitor. Because MAO inhibitors have been widely applied for treating depression, xanthoangelol (1) and 4-hydroxyderricin (3) were suggested as candidates for developing combined antidepressant drug. Additionally, cynaroside (43, luteolin-7-O- β -D-glucopyranoside) showed strong inhibitory effect against DBH (Kim et al. 2013).

Anti-viral activity

Park et al. reported that xanthoangelol F (2), 4-hydroxyderricin (3), xanthoangelol B (5), xanthoangelol G (6), xanthoangelol D (9), and xanthokeistal A (28) have inhibitory effects against hydrolysis of influenza virus neuraminidase (NA) which has been an effective target to develop drugs treating patients infected with influenza

virus. All active compounds showed reversible nonselective characteristics and xanthoangelol D (**9**) was the most potent inhibitor. Furthermore, it was suggested that the influenza virus NA inhibitory activities of the alkylated chalcones are associated with their alkyl groups (Park et al. 2011). Because enzymatic activities of a chymotrypsin-like protease (3CL^{pro}) and a papain-like protease (PL^{pro}) are essential for a viral life cycle of severe acute respiratory syndrome coronavirus (SARS-CoV), the two viral cysteine proteases are attractive targets for development of anti-SARS-CoV drugs. Xanthoangelol (**1**), xanthoangelol F (**2**), 4-hydroxyderricin (**3**), isobavachalcone (**4**), xanthoangelol B (**5**), xanthoangelol G (**6**), xanthoangelol D (**9**), xanthoangelol E (**10**), and xanthokeistal A (**28**) showed competitive inhibitory characteristics to 3CL^{pro} and noncompetitive ones to PL^{pro}, except for mixed inhibition manner of isobavachalcone (**4**) against PL^{pro}. Especially, xanthoangelol E (**10**) exhibited the most potent inhibitory activity, and docking simulation of xanthoangelol E (**10**) to both protease revealed that its hydroperoxy group plays a role in enzyme binding by forming hydrogen bonding (Park et al. 2015).

Osteogenesis promotive activity

Angelichalcone (**31**) was presented as an osteogenesis promotive agent in a patent application (Ohnogi et al. 2004).

Whitening activity

Xanthoangelol (**1**), 4-hydroxyderricin (**3**), xanthoangelol H (**12**), deoxyxanthoangelol H (**13**), and deoxydihydroxanthoangelol H (**41**) exhibited inhibitory effects against melanin formation in B16 melanoma cells with low cytotoxicity (Arung et al. 2012).

Discussion

As presented in this review, the pharmacological studies on *A. keiskei* and its putative active compounds support a number of biological activities that can potentially impact human health. In spite of these reported biological activities and a number of filed patent applications on this plant, *A. keiskei* has not been developed as a pharmaceutical agent and is currently available on the market as health drinks. This review presented a summary of the studies published to date on this promising plant. Although the majority of these studies are mechanistic, but they can be used as a solid platform to design and conduct clinical studies.

Acknowledgements This work is supported by Ewha Womans University (collaborative research grant for Ewha-UC Irvine).

Compliance with ethical standards

Conflict of interest The co-authors have declared no conflict of interest.

References

- Akihisa T, Tokuda H, Ukiya M, Iizuka M, Schneider S, Ogasawara K, Mukainaka T, Iwatsuki K, Suzuki T, Nishino H (2003) Chalcones, coumarins, and flavanones from the exudate of *Angelica keiskei* and their chemopreventive effects. *Cancer Lett* 201:133–137
- Akihisa T, Tokuda H, Hasegawa D, Ukiya M, Kimura Y, Enjo F, Suzuki T, Nishino H (2006) Chalcones and other compounds from the exudates of *Angelica keiskei* and their cancer chemopreventive effects. *J Nat Prod* 69:38–42
- Akihisa T, Yasukawa T, Ukiya M, Hasegawa D (2007) Anti-inflammatory agents containing natural products from *Angelica keiskei*. (Nihon University, Japan). Application: JP
- Akihisa T, Kikuchi T, Nagai H, Ishii K, Tabata K, Suzuki T (2011) 4-Hydroxyderricin from *Angelica keiskei* roots induces caspase-dependent apoptotic cell death in HL60 human leukemia cells. *J Oleo Sci* 60:71–77
- Aoki N, Ohta S (2010) Ashitabaol A, a new antioxidative sesquiterpenoid from seeds of *Angelica keiskei*. *Tetrahedron Lett* 51:3449–3450
- Aoki N, Muko M, Ohta E, Ohta S (2008) C-Geranylated chalcones from the stems of *Angelica keiskei* with superoxide-scavenging activity. *J Nat Prod* 71:1308–1310
- Arung ET, Furuta S, Sugamoto K, Shimizu K, Ishikawa H, Matsushita Y-I, Kondo R (2012) The inhibitory effects of representative chalcones contained in *Angelica keiskei* on melanin biosynthesis in B16 melanoma cells. *Nat Prod Commun* 7:1007–1010
- Baba K, Kido T, Yoneda Y, Taniguchi M, Kozawa M (1990a) Chemical components of *Angelica keiskei* Koidzumi. (V). Components of the fruits, and comparison of coumarins and chalcones in the fruits, roots and the leaves. *Shoyakugaku Zasshi* 44:235–239
- Baba K, Nakata K, Taniguchi M, Kido T, Kozawa M (1990b) Chemical components of *Angelica keiskei*. Part 7. Chalcones from *Angelica keiskei*. *Phytochemistry* 29:3907–3910
- Baba K, Taniguchi M, Nakata K (1998) Studies on *Angelica keiskei* “Ashitaba”. *Foods Food Ingrid J Jpn* 178:52–60
- Bao G (2014) *Angelica keiskei*-containing nutritional and health-protecting flour and its preparation method (Qingyang county nanyang rice industry Co., Ltd, Peop. Rep. China). Application: CN
- Battenberg OA, Yang Y, Verhelst SHL, Sieber SA (2013) Target profiling of 4-hydroxyderricin in *S. aureus* reveals seryl-tRNA synthetase binding and inhibition by covalent modification. *Mol Biosyst* 9:343–351
- Chang HR, Lee HJ, Ryu J-H (2014) Chalcones from *Angelica keiskei* attenuate the inflammatory responses by suppressing nuclear translocation of NF-κB. *J Med Food* 17:1306–1313
- Enoki T, Ohnogi H, Nagamine K, Kudo Y, Sugiyama K, Tanabe M, Kobayashi E, Sagawa H, Kato I (2007) Antidiabetic activities of chalcones isolated from a Japanese herb, *Angelica keiskei*. *J Agric Food Chem* 55:6013–6017
- Fujita T, Sakuma S, Sumiya T, Nishida H, Fujimoto Y, Baba K, Kozawa M (1992) The effects of xanthoangelol E on arachidonic

- acid metabolism in the gastric antral mucosa and platelet of the rabbit. *Res Commun Chem Pathol Pharmacol* 77:227–240
- Fukuo Y, Ogasawara K, Satoh K (2005) Antimicrobial agent/ Antimicrobial agent against *Helicobacter pylori* comprising a material derived from *Angelica keiskei* (Japan Bio Science Laboratory Co., Ltd., Japan). Application: US
- Guo X-Q, Wu J-H, Liu C, Li J-Q, Wang Z-W, Yang K-F (2013) Comparison of antioxidant content and oxidation resistance of stems and leaves from *Angelica keiskei* Koidzumi. *Shipin Yu Fajiao Gongye* 39:122–127
- Hata K, Kozawa M (1961) Pharmacognostical studies on umbelliferous plants. XVIII. Constituents of the roots of *Angelica keiskei*. *Yakugaku Zasshi* 81:1647–1649
- Higa Y (1985) Skin-whitening cosmetic containing *Angelica keiskei* Koidzumi extract and having tyrosinase inhibiting effect (Sansei Pharmaceutical Co., Ltd., Japan). Application: JP
- Imai K, Imai M (2008) Encyclopedia of wild herbs and vegetables. Green Home, Seoul
- Inamori Y, Baba K, Tsujibo H, Taniguchi M, Nakata K, Kozawa M (1991) Chemical components of *Angelica keiskei*. VI. Antibacterial activity of two chalcones, xanthoangelol and 4-hydroxyderricin, isolated from the root of *Angelica keiskei* Koidzumi. *Chem Pharm Bull* 39:1604–1605
- Jeong Y-J, Kang KJ (2011) Effect of *Angelica keiskei* extract on apoptosis of MDA-MB-231 human breast cancer cells. *Han'guk Sikp'um Yongyang Kwahak Hoechi* 40:1654–1661
- Kang M-H, Park YK, Kim H-Y, Kim TS (2004) Green vegetable drink consumption protects peripheral lymphocytes DNA damage in Korean smokers. *BioFactors* 22:245–247
- Kawabata K-I, Sawada K, Ikeda K, Fukuda I, Kawasaki K, Yamamoto N, Ashida H (2011) Prenylated chalcones 4-hydroxyderricin and xanthoangelol stimulate glucose uptake in skeletal muscle cells by inducing GLUT4 translocation. *Mol Nutr Food Res* 55:467–475
- Kil Y-S, Choi S-K, Lee Y-S, Jafari M, Seo E-K (2015a) Chalcones from *Angelica keiskei*: evaluation of their heat shock protein inducing activities. *J Nat Prod* 78:2481–2487
- Kil Y-S, Nam J-W, Lee J, Seo EK (2015b) Separation of two major chalcones from *Angelica keiskei* by high-speed counter-current chromatography. *Arch Pharmacol Res* 38:1506–1511
- Kil Y-S, Kwon J, Lee D, Seo EK (2016) Three new chalcones from the aerial parts of *Angelica keiskei*. *Helv Chim Acta* 99:393–397
- Kim S-J, Cho J-Y, Wee J-H, Jang M-Y, Kim C, Rim Y-S, Shin S-C, Ma S-J, Moon J-H, Park K-H (2005) Isolation and characterization of antioxidative compounds from the aerial parts of *Angelica keiskei*. *Food Sci Biotechnol* 14:58–63
- Kim TS, Shim JS, Kim SD, Yeo IH (2006) Health-aid food, drink and food composition having hepato-protective effect, which comprises *Angelica keiskei* root extract and hydroxyderricin and xanthoangelol isolated therefrom (Pulmuone Co., Ltd., S. Korea). Application: KR
- Kim E, Choi J, Yeo I (2012) The effects of *Angelica keiskei* Koidz on the expression of antioxidant enzymes related to lipid profiles in rats fed a high fat diet. *Nutr Res Pract* 6:9–15
- Kim JH, Son YK, Kim GH, Hwang KH (2013) Xanthoangelol and 4-hydroxyderricin are the major active principles of the inhibitory activities against monoamine oxidases on *Angelica keiskei* K. *Biomol Ther* 21:234–240
- Kim DW, Curtis-Long MJ, Yuk HJ, Wang Y, Song YH, Jeong SH, Park KH (2014) Quantitative analysis of phenolic metabolites from different parts of *Angelica keiskei* by HPLC-ESI MS/MS and their xanthine oxidase inhibition. *Food Chem* 153:20–27
- Kimura Y, Baba K (2003) Antitumor and antimetastatic activities of *Angelica keiskei* roots, part 1: isolation of an active substance, xanthoangelol. *Int J Cancer* 106:429–437
- Kimura Y, Taniguchi M, Baba K (2004) Antitumor and antimetastatic activities of 4-hydroxyderricin isolated from *Angelica keiskei* roots. *Planta Med* 70:211–219
- Kimura Y, Sumiyoshi M, Baba K (2008) Anti-tumor actions of major component 3'-O-acetylhamaudol of *Angelica japonica* roots through dual actions, anti-angiogenesis and intestinal intraepithelial lymphocyte activation. *Cancer Lett* 265:84–97
- Kishiro S, Nunomura S, Nagai H, Akihisa T, Ra C (2008) Selinidin suppresses IgE-mediated mast cell activation by inhibiting multiple steps of FcεRI signaling. *Biol Pharm Bull* 31:442–448
- Kozawa M, Morita N, Baba K, Hata K (1977) The structure of xanthoangelol, a new chalcone from the roots of *Angelica keiskei* Koidzumi (Umbelliferae). *Chem Pharm Bull* 25:515–516
- Kozawa M, Morita N, Baba K, Hata K (1978) Chemical components of the roots of *Angelica keiskei* Koidzumi. III. The structure of a new dihydrofurocoumarin. *Yakugaku Zasshi* 98:636–638
- Ku CS, Kim MJ, Jung YJ, Jang DI (2014) Cosmetic compositions containing *Angelica keiskei* extracts (COTDE Co., Ltd., S. Korea). Application: KR
- Kusaba N, Kamiya T, Ikeguchi M, Ono H, Takagaki K, Hayashi M, Ogasawara T, Sugawa-Katayama Y (2008) Effects of AOJIRU drink powder of *Angelica keiskei* leaf containing indigestible dextrin on defecation in volunteers with constipation. *Jpn Pharmacol Ther* 36:1159–1165
- Lee HJ, Choi TW, Kim HJ, Nam D, Jung SH, Lee EH, Lee HJ, Shin EM, Jang H-J, Ahn KS, Shim BS, Choi S-H, Kim S-H, Sethi G, Ahn KS (2010) Anti-inflammatory activity of *Angelica keiskei* through suppression of mitogen-activated protein kinases and nuclear factor-κB activation pathways. *J Med Food* 13:691–699
- Lee MS, Jung MG, Choi GB, Kim TS, Yeo IH, Nam SU (2012a) *Angelica keiskei* extracts for preventing and treating obesity (Pulmuone Holdings Co., Ltd., S. Korea). Application: KR
- Lee MS, Kim BG, Choi GB, Kim TS, Yeo IH, Nam SU (2012b) Skin whitening composition containing *Angelica keiskei* extracts (Pulmuone Holdings Co., Ltd., S. Korea). Application: KR
- Lee MS, Lee EJ, Choi GB, Kim TS, Yeo IH, Nam SU (2013) *Angelica keiskei* extracts for preventing and alleviating diabetes mellitus (Pulmuone Holdings Co., Ltd., S. Korea). Application: KR
- Lee MS, Choi GB, Kim TS, Yeo IH, Nam SU (2014) *Angelica keiskei* root extracts for inducing apoptosis of prostate cancer cells (Pulmuone Co., Ltd., S. Korea). Application: KR
- Li J-L, Gao L-X, Meng F-W, Tang C-L, Zhang R-J, Li J-Y, Luo C, Li J, Zhao W-M (2015) PTP1B inhibitors from stems of *Angelica keiskei* (Ashitaba). *Bioorg Med Chem Lett* 25:2028–2032
- Liu X, Wang Z, Sun J, Wan Y, Rong Y (2014) Coix seed-*Angelica keiskei* nutritional health-care biscuits and production method thereof (Shanghai Jiao Tong University, Peop. Rep. China). Application: CN
- Luo L, Wang R, Wang X, Ma Z, Li N (2012) Compounds from *Angelica keiskei* with NQO1 induction, DPPH· scavenging and α-glucosidase inhibitory activities. *Food Chem* 131:992–998
- Matsuura M, Kimura Y, Nakata K, Baba K, Okuda H (2001) Artery relaxation by chalcones isolated from the roots of *Angelica keiskei*. *Planta Med* 67:230–235
- Motani K, Tabata K, Kimura Y, Okano S, Shibata Y, Abiko Y, Nagai H, Akihisa T, Suzuki T (2008) Proteomic analysis of apoptosis induced by xanthoangelol, a major constituent of *Angelica keiskei*, in neuroblastoma. *Biol Pharm Bull* 31:618–626
- Murakami S, Kijima H, Isobe Y, Muramatsu M, Aihara H, Otomo S, Baba K, Kozawa M (1990) Inhibition of gastric H⁺, K⁺-ATPase by chalcone derivatives, xanthoangelol and 4-hydroxyderricin, from *Angelica keiskei* Koidzumi. *J Pharm Pharmacol* 42:723–726

- Murakami S, Kotomo S, Ozawa M, Baba K (1992) Novel chalcone derivative as antiulcer agent (Taisho Pharmaceutical Co., Ltd., Japan). Application: JP
- Nakata K, Baba K (2001) Histamine release-inhibiting activity of *Angelica keiskei*. *Nat Med* 55:32–34
- Nakata K, Katsumata H, Taniguchi M, Kita S, Baba K (1997) Comparison of the characteristics and the constituents of *Angelica keiskei* KOIDZUMI growing in Hachijo and Oshima. *Nat Med* 51:532–536
- Nakata K, Taniguchi M, Baba K (1999) Three chalcones from *Angelica keiskei*. *Nat Med* 53:329–332
- Nishimura R, Tabata K, Arakawa M, Ito Y, Kimura Y, Akihisa T, Nagai H, Sakuma A, Kohno H, Suzuki T (2007) Isobavachalcone, a chalcone constituent of *Angelica keiskei*, induces apoptosis in neuroblastoma. *Biol Pharm Bull* 30:1878–1883
- Noh H-M, Ahn E-M, Yun J-M, Cho B-L, Paek Y-J (2015) *Angelica keiskei* Koidzumi extracts improve some markers of liver function in habitual alcohol drinkers: a randomized double-blind clinical trial. *J Med Food* 18:166–172
- Ogawa H, Nakashima S, Baba K (2003) Effects of dietary *Angelica keiskei* on lipid metabolism in stroke-prone spontaneously hypertensive rats. *Clin Exp Pharmacol Physiol* 30:284–288
- Ogawa H, Nakamura R, Baba K (2005a) Beneficial effect of laserpitin, a coumarin compound from *Angelica keiskei*, on lipid metabolism in stroke-prone spontaneously hypertensive rats. *Clin Exp Pharmacol Physiol* 32:1104–1109
- Ogawa H, Ohno M, Baba K (2005b) Hypotensive and lipid regulatory actions of 4-hydroxyderricin, a chalcone from *Angelica keiskei*, in stroke-prone spontaneously hypertensive rats. *Clin Exp Pharmacol Physiol* 31:19–23
- Oh SR, Kim S-J, Kim DH, Ryu JH, Ahn E-M, Jung JW (2013) *Angelica keiskei* ameliorates scopolamine-induced memory impairments in mice. *Biol Pharm Bull* 36:82–88
- Ohkura N, Nakakuki Y, Taniguchi M, Kanai S, Nakayama A, Ohnishi K, Sakata T, Nohira T, Matsuda J, Baba K, Atsumi G-I (2011) Xanthoangelols isolated from *Angelica keiskei* inhibit inflammatory-induced plasminogen activator inhibitor 1 (PAI-1) production. *BioFactors* 37:455–461
- Ohnogi H, Shiraga M, Sagawa H, Kato I (2002) Nerve growth factor enhancers for treatment of disease containing coumarin and/or chroman derivatives (Takara Bio Inc., Japan). Application: WO
- Ohnogi H, Sugiyama K, Muraki N, Sagawa H, Kato I (2004) Plant components for promoting osteogenesis (Takara Bio Inc., Japan). Application: WO
- Ohnogi H, Enoki T, Hino F, Kato I (2007) Antidiabetic effect and safety of long-term ingestion of Ashitaba (*Angelica keiskei*) powder containing Chalcone (4HD) on borderline mild hyperglycemia. *Jpn Pharmacol Ther* 35:647–660
- Ohnogi H, Hayami S, Kudo Y, Deguchi S, Mizutani S, Enoki T, Tanimura Y, Aoi W, Naito Y, Kato I, Yoshikawa T (2012a) *Angelica keiskei* extract improves insulin resistance and hypertriglyceridemia in rats fed a high-fructose drink. *Biosci Biotechnol Biochem* 76:928–932
- Ohnogi H, Kudo Y, Tahara K, Sugiyama K, Enoki T, Hayami S, Sagawa H, Tanimura Y, Aoi W, Naito Y, Kato I, Yoshikawa T (2012b) Six new chalcones from *Angelica keiskei* inducing adiponectin production in 3T3-L1 adipocytes. *Biosci Biotechnol Biochem* 76:961–966
- Oi J, Meyer FG, Meyer FG, Walker EH (1965) *Flora of Japan*: in English: combined, much revised and extended translation/by the author of his *Flora of Japan* (1953) and *Flora of Japan*, Pteridophyta (1957); edited by Frederick G. Meyer and Egbert H. Walker. Washington, Smithsonian Institution
- Okuyama T, Takata M, Takayasu J, Hasegawa T, Tokuda H, Nishino A, Nishino H, Iwashima A (1991) Anti-tumor-promotion by principles obtained from *Angelica keiskei*. *Planta Med* 57:242–246
- Onogi H, Sugiyama K, Muraki N, Enoki T, Sagawa H, Kato I (2004) Chalcone derivatives from *Angelica* and pharmacological activities thereof (Takara Bio Inc., Japan). Application: WO
- Park J-C (2013) The medicinal herbs, teas, and alcoholic beverages in Donguibogam, proven by patents. Pureun-Hyeongbok Publisher, Goyang
- Park JC, Cho YS, Park SK, Park JR, Chun SS, Ok KD, Choi JW (1995) Isolation of flavone-7-O-glycosides from the aerial parts of *Angelica keiskei* and antihyperlipidemic effects. *Saengyak Hakhoechi* 26:337–343
- Park JC, Yu YB, Lee JH, Choi MR, Ok KD (1996) Chemical components from the aerial parts of *Angelica keiskei*. *Saengyak Hakhoechi* 27:80–82
- Park J-C, Park J-R, Chung S-K, Yu Y-B, Ha J-O, Park K-Y (1997a) Antimutagenic activity of the methanol extract and compounds of *Angelica keiskei* in the Salmonella assay system. *Saengyak Hakhoechi* 28:80–83
- Park J-R, Park J-C, Choi S-H (1997b) Screening and characterization of anticholesterogenic substances from food plant extracts. *Han'guk Sikk'um Yongyang Kwahak Hoechi* 26:236–241
- Park JC, Park JG, Kim HJ, Hur JM, Lee JH, Sung NJ, Chung SK, Choi JW (2002) Effects of extract from *Angelica keiskei* and its component, cynaroside, on the hepatic bromobenzene-metabolizing enzyme system in rats. *Phytother Res* 16:S24–S27
- Park J-Y, Jeong HJ, Kim YM, Park S-J, Rho M-C, Park KH, Ryu YB, Lee WS (2011) Characteristic of alkylated chalcones from *Angelica keiskei* on influenza virus neuraminidase inhibition. *Bioorg Med Chem Lett* 21:5602–5604
- Park JY, Ko Jin A, Kim Dae W, Kim Young M, Kwon HJ, Jeong Hyung J, Kim Cha Y, Park Ki H, Lee Woo S, Ryu Young B (2015) Chalcones isolated from *Angelica keiskei* inhibit cysteine proteases of SARS-CoV. *J Enzym Inhib Med Chem*: 1–8
- Qin LQ, Luo SY, Zhan ZH, Liu XX, Wang K (2014) Determination of antioxidant compounds from leaf and stem of *Angelica keiskei* by gas chromatography-mass spectrometry. *Asian J Chem* 26:5097–5099
- Saeki Y, Asada S, Kikuchi S, Itoh S (2013) Deodorant composition containing extracts of *Angelica keiskei* for prevention of mouth odor (Lotte Co., Ltd., Japan). Application: WO; Chemical Indexing Equivalent to 158:544360 (JP)
- Sakaguchi Y, Sakaguchi M, Sakaguchi T (1996) Hair growth promoter comprising *Angelica keiskei* extract, and preparation method thereof (Japan). Application: JP
- Shimizu E, Hayashi A, Takahashi R, Aoyagi Y, Murakami T, Kimoto K (1999) Effects of angiotensin 1-converting enzyme inhibitor from Ashitaba (*Angelica keiskei*) on blood pressure of spontaneously hypertensive rats. *J Nutr Sci Vitaminol* 45:375–383
- Shin JE, Choi EJ, Jin Q, Jin H-G, Woo E-R (2011) Chalcones isolated from *Angelica keiskei* and their inhibition of IL-6 production in TNF- α -stimulated MG-63 cell. *Arch Pharmacol Res* 34:437–442
- Shin H-J, Shon D-H, Youn H-S (2013) Isobavachalcone suppresses expression of inducible nitric oxide synthase induced by Toll-like receptor agonists. *Int Immunopharmacol* 15:38–41
- Son DJ, Park YO, Yu C, Lee SE, Park YH (2014) Bioassay-guided isolation and identification of anti-platelet-active compounds from the root of Ashitaba (*Angelica keiskei* Koidz.). *Nat Prod Res* 28:2312–2316
- Su L (2014) A method for preparing *Angelica keiskei* wine (Nanjing Zelang Agriculture Development Co., Ltd., Peop. Rep. China). Application: CN
- Sugamoto K, Matsushita Y-I, Matsui T, Kurogi C, Park JC (2008) Isolation and analysis components of Ashitaba (*Angelica keiskei* koidzumi) leaves growing in Miyazaki and Korea. *Mem Fac Eng Miyazaki Univ* 37:75–80

- Sugamoto K, Matsushita Y-I, Matsui T, Hikita H, Park JC (2009) Isolation and analysis components of Ashitaba (*Angelica keiskei* Koidzumi) roots, stems and leaves. Mem Fac Eng Miyazaki Univ 38:91–96
- Sugamoto K, Matsusita Y-I, Matsui K, Kurogi C, Matsui T (2011) Synthesis and antibacterial activity of chalcones bearing prenyl or geranyl groups from *Angelica keiskei*. Tetrahedron 67:5346–5359
- Sugii M, Ohkita M, Taniguchi M, Baba K, Kawai Y, Tahara C, Takaoka M, Matsumura Y (2005) Xanthoangelol D isolated from the roots of *Angelica keiskei* inhibits endothelin-1 production through the suppression of nuclear factor- κ B. Biol Pharm Bull 28:607–610
- Tabata K, Motani K, Takayanagi N, Nishimura R, Asami S, Kimura Y, Ukiya M, Hasegawa D, Akihisa T, Suzuki T (2005) Xanthoangelol, a major chalcone constituent of *Angelica keiskei*, induces apoptosis in neuroblastoma and leukemia cells. Biol Pharm Bull 28:1404–1407
- Takaoka S, Hibasami H, Ogasawara K, Imai N (2008) Chalcones from *Angelica keiskei* induced apoptosis in stomach cancer cells. J Herbs, Spices Med Plants 14:166–174
- Yasuda M, Kawabata K, Miyashita M, Okumura M, Yamamoto N, Takahashi M, Ashida H, Ohigashi H (2014) Inhibitory effects of 4-hydroxyderricin and xanthoangelol on lipopolysaccharide-induced inflammatory responses in RAW264 macrophages. J Agric Food Chem 62:462–467
- Zhang T, Sawada K, Yamamoto N, Ashida H (2013) 4-Hydroxyderricin and xanthoangelol from Ashitaba (*Angelica keiskei*) suppress differentiation of preadipocytes to adipocytes via AMPK and MAPK pathways. Mol Nutr Food Res 57:1729–1740
- Zhang T, Yamashita Y, Yasuda M, Yamamoto N, Ashida H (2015) Ashitaba (*Angelica keiskei*) extract prevents adiposity in high-fat diet-fed C57BL/6 mice. Food Funct 6:134–145
- Zhou X, Liang C, Xu Q, Zhang X, Liang X, Huang Q, Su X (2012) Chemical constituents of *Angelica keiskei*. Zhongguo Shiyang Fangjixue Zazhi 18:103–105